

RSV can cause:

- Fever
- Runny Nose
- Sore Throat
- Ear Infection
- Cough
- Croup
- Bronchiolitis
(infection of the lungs)
- Pneumonia
- Wheezing



We need your help!

Participants must:

- Be less than 2 years old.
- Have signs and symptoms of a respiratory illness in the last 5 days.

Principal Investigators

Janet Englund, MD
Seattle Children's
Infectious Disease

Eileen Klein, MD
Seattle Children's
Emergency Department

What is this research about?

- Respiratory Syncytial Virus (RSV) is a germ.
- RSV can make infants and young children very sick.
- Researchers want to know:
 - What percentage of children with RSV is hospitalized?
 - How much RSV is in the nose during infection?

What would I do?

- Participation lasts about 2 weeks. There would be 1 visit and 1 follow-up phone call/email.
- Participants are asked to:
 - Give a nasal sample by nasal swab. We would test the swab for RSV.
 - Allow researchers to review their child's medical record about their child's illness.

What else should I know?

- The study visit can be at your clinic, Seattle Children's, or your home. The visit takes 15-30 minutes.
- You would be paid for your time in the study
- All services and visits are at no charge to you.
- Participation is voluntary.

Where can I get more information?

Ask your clinic if a Seattle Children's research team member is available or:

Phone: (206) 884-1100

Email: idresearch@seattlechildrens.org